

Modelling of long-term future climate change with application to the problem of permanent nuclear waste storage in Germany

Christine Kaufhold, Dr. Andrey Ganopolski

Earth System Analysis Department
Potsdam Institute for Climate Impact Research
March 17, 2023




BUNDESGESELLSCHAFT
FÜR ENDLAGERUNG



POTSDAM-INSTITUT FÜR
KLIMAFOLGENFORSCHUNG

GRS URS2023



Our past climate...

- Climate has been changing for last Myr via Milankovitch cycles
- Connection between maximum summer insolation at 65°N and CO₂ for glacial inception




Figure 1: Data from Quinn et al. 1991/Liseki & Raymo 2005. Generated by Rohde 2006.





Figure 2: Critical CO₂-insolation relation for glacial inception. From Ganopolski 2016, *Nature*.

Our future climate...

- Available fossil fuel carbon reserves have the capacity to impact the climate hundreds of thousands of years into the future
- Large uncertainties exist in long-term future climate evolution since we cannot accurately predict anthropogenic CO₂ emissions even during this century
 - RCP-SSP scenarios with extensions
- Anthropogenic CO₂ emissions can change future glacial cycles

Figure 3: Taken from IPCC's AR6.



How does this affect the management of nuclear waste?

- Previous glaciation events reached Germany
 - many short-term and long-term consequences
 - precipitation, temperature, subterranean stress, surface denudation, and permafrost/taliks
- Why does this matter for repository health?
 - some radioisotopes have long half lives
 - waste must be stored for a period of 1 million years (EndSiAnfV § 3)
 - **we must consider future climate for deep geological repositories**



The REDUKLIM project

Research Field 4:

Preliminary safety investigation

Research Cluster:

Uncertainties and Robustness with regard to the Safety of a repository for high-level radioactive waste (URS)

Topic 4:

Physics-based scenario modelling and impact models

Project:

Reduction of scenario uncertainties through Climate models (REDUKLIM)




Figure 4: Structure of the URS cluster from BGE

The goals of REDUKLIM

The task:

- assess future climate for next 100 kyr and 1 Myr
- link climate development to groundwater processes (GRS)
- identify and quantify uncertainties via our projections
- **provide additional confidence in site selection**

Our tools at hand:

1. distributed density-driven flow model d^3f_{++} (GRS)
2. Earth system model of intermediate complexity CLIMBER-X
3. reduced complexity model by Talento & Ganopolski 2021





Figure 5: Areas in consideration. Colours symbolize the different host rocks. Taken from BGE.



- Fully coupled EMIC
- Climate components have comparable complexity and grid ($5^\circ \times 5^\circ$)
- More options for the ice sheet model
 - e.g., variable domain
- Best suited for long timescales
 - does not resolve weather, inter-annual variability, diurnal cycle, etc.
 - we will use until ~ 100 kyr AP
- Model validation in Willeit et al. 2022



A reduced complexity model

3 coupled, nonlinear equations concerning mechanisms relevant for the climate–icesheet–carbon cycle system on very long timescales (> 10 kyr)





Figure 6: Talento & Ganopolski 2021

Timeline of PhD and REDUKLIM




Package milestone #1

Working Package 1.1:

Construction and evaluation of climate models

1. simulate the last glacial cycle
2. simulate the next 100,000 yrs for different scenarios
3. for 100,000 yrs and beyond, the reduced complexity model is used
4. scenarios developed for the next million years in northern Germany/Alpine region




Milestone 1.11 (month 12):

Evaluation of future climate scenarios for Germany based on results of CLIMBER-X

Paper 1: Glacial cycles

- **To simulate future glacial cycles, we must be able to simulate the last one**
 - glacial inception, LGM, and deglaciation
- Major objective as model performance can be tested against paleodata
 - sediment core, ice core, planktonic data
 - reconstructions (GLAC-1D, ICE-6G_C)
 - sea level reconstructions
 - PMIP4 model ensemble results
- Currently status:
 - model tuning
 - identifying model biases
 - drafting paper (e.g., introduction, methods)



Paper 1: Transient simulation

"a serious impediment... is that no modern model of the coupled climate system has ever been shown to naturally produce such oscillatory behavior under glacial climate conditions"


- **Peltier 2014, Geophysical Research Letters**

"this deficiency is related to both the computational expense which prevents models from being run for the longer time periods..."



- **Malmierca-Vallet 2022, EGU sphere**

"a full transient glacial cycle is currently computationally unfeasible as it requires a too-large amount of computation time"


- **Scherrenberg 2023, Climate of the Past**




Paper 1: Preliminary results





ICE-6G_C



GLAC-1D



CLIMBER-X



Transient simulations and model-data comparison of the last glacial cycle using a coupled climate-ice sheet model

Christine Kaufhold^{1,2}, Matteo Willeit¹, and Andrey Ganopolski¹

¹Earth System Analysis, Potsdam Institute for Climate Impact Research, Potsdam, Germany

²Institute of Physics and Astronomy, University of Potsdam, Potsdam, Germany

Correspondence: Christine Kaufhold (kaufhold@pik-potsdam.de)

Abstract. TEXT

1 Introduction

The last few million years in Earth's history saw generated a large variability in our climate (Lisiecki and Raymo, 2005). As a result, there are still many unknowns as global proxy data on seasonal to millennial timescales from sources ranging from


- 5 ice cores to marine sediments has provided poor constraints on paleoclimatic conditions. Yet one such exception is the last glacial maximum (LGM) of the Pleistocene epoch, which is generally suggested to have occurred sometime between 24.5 to 17 kyr BP (Clark et al., 2009). Ice sheets at this time are generally well constrained by present-day observations in addition to terrestrial (tree rings, sediments), ice (e.g., stacked $\delta^{18}\text{O}$ data in cores) and marine indicators (e.g., lake sediments, ice-rafted debris). LGM can be distinctly recognized in paleoclimate records by a large volume of Northern Hemisphere (NH) ice sheets and correspondingly low global eustatic sea level (Fairbanks, 1989; Yokoyama et al., 2000; Waelbroeck et al., 2002; Pelicci
- 10

Package milestone #2

Working Package 1.2:

Scenarios, uncertainties, robustness

1. examine uncertainty/consequences of a wide range of cumulative CO₂ scenarios
2. examine how glacial inception depends on parameters of the carbon component
3. explore uncertainties in climate sensitivity and ice sheet parameterization




Milestone 1.21 (month 24):


Assessment of the robustness and uncertainties of future climate scenarios for Germany

Paper 2: Future scenarios


- Climate scenarios offer a spectrum of possible model outcomes
- Simulate long-term scenarios for a broad range of cumulative CO₂ emissions
- **How do cumulative CO₂ emissions affect timing of the onset of next glaciation?**
- Prescribe CO₂:
 - interactive open carbon cycle
 - reduced complexity model
 - Lord et al. 2016
- range of possible climates over Europe (temperature, precipitation, sea level, etc.)



Paper 2: Preliminary results




CLIMBER X
Earth System Model



Ganopolski 2016
Critical CO₂ CLIMBER-2:



$$smx65_{cr} = -77 \log \left(\frac{CO_2}{280} \right) + 466$$



Paper 2: Validation

- Lord 2016 created an emulator for CO₂ emissions pulses
- Volcanic outgassing is tuned to balance silicate weathering O(20,000 yrs) over one glacial cycle
- Further investigation on this relation is required

Lord 2016:



Collaborations and coauthorships

- Successful Climate workshop organized by BGE in February 2023
- Lots of interest for potential collaborations:
 - **Erosion project** with Todd Ehlers
 - **QUASI project** with Anders Damsgaard
- At least 1 (review) coauthorship planned with Andrey Ganopolski on deep-future climate modelling
 - what's been done/general literature review
 - certainties and uncertainties in the climate
 - Anthropocene-Pleistocene II transition
 - potential late Pleistocene II transition?

Agenda Climate Workshop		BGE BUNDESGESELLSCHAFT FÜR ENDLAGERUNG
		06.02.2023 09:00 Uhr bis 15:00 Uhr BGE mbH Peme (Slope Meeting) Ansprechpartner: Marc Wengler
Participants:		
- REDUKLIM: Judith Flügge (GRS), Marc Johnen (GRS), Jens Wolf (GRS), Andrey Ganopolski (PIK), Christine Kaufhold (PIK)		
- Subrosion ewG: Judith Flügge (GRS), Christine Fahrholz (GRS)		
- QUASI: Jan Piotrowski (Uni Aarhus), Anders Damsgaard (Uni Aarhus), Jutta Wissemann (Uni Hannover), Runa Fäber (Uni Hannover)		
- Suchtiefe: Jörg Lang (BGR)		
- Zyklische Vergleitscherungen: Tobias Baumann (smartTectonics)		
- Erosion: Todd Ehlers (Uni Tübingen), Christoph Glötzbach (Uni Tübingen)		
- BGE: Anne Bartelzko, Wolfram Rühsek, Eva-Maria Hoyer, Paulina Müller, Florian Schöne, Marc Wengler, Nils-Peter Nilus, Julia Riemacker-Burschli, Nadine Schöne		
TOP 1	09:00 – 09:05	Welcome address (BGE)
TOP 2	09:05 – 09:15	Introduction of participants
TOP 3	09:15 – 12:15	Presentations of the projects (15 min presentation + 15 min Q&A)
	TOP 3.1	09:15 – 09:45 REDUKLIM
	TOP 3.2	09:45 – 10:15 Subrosion ewG
	TOP 3.3	10:15 – 10:45 QUASI
	TOP 3.4	10:45 – 11:15 Suchtiefe
	TOP 3.5	11:15 – 11:45 Zyklische Vergleitscherungen
	TOP 3.6	11:45 – 12:15 Erosion
TOP 4	12:15 – 13:15	Break
TOP 5	13:15 – 13:45	Presentation of the FEP methodology (BGE)
TOP 6	13:45 – 15:00	Discussion

Strategy & outlook

- Learnt CLIMBER-X over the course of the last months and made good progress
- Clear aim in terms of what must be done & the general timeline
- Few ideas to navigate bumps and prepare for success

Funded by:



Project Information:

Ungewissheiten und Robustheit mit Blick auf die Sicherheit eines Endlagers für hochradioaktive Abfälle (bge.de)

Contact:

kaufhold@pik-potsdam.de